

BORN nutrition

Provides Musculoskeletal Support
Provides Bone Support | Increases Skeletal Strength and Helps
Maintain Bone Density



Strontium

What is Strontium?

Strontium is a mineral that is similar physical and chemical properties to calcium. Research has shown strontium provides bone support through its ability to increase the formation of osteoblasts (cells that build up bone) and slow down the formation of osteoclasts (cells that break down bone tissue) helping to maintain healthy bone density. As a result, strontium imparts a balancing effect on the osteoblastic-osteoclastic ratio and boosts skeletal strength via two different mechanisms. Strontium's ability to strengthen both the bone matrix and skeletal density makes it an important part of any bone-building regimen.

Overview

Bone is the rigid structure that makes up our skeleton, but is actually an active, living organ that is constantly being built up and broken down by osteoblasts and osteoclasts. Aging, genetic predisposition, lifestyle factors, and co-morbidities can alter the balance between breakdown and building of new bone structure, bone strength, and flexibility. Mineral balance is one nutritional factor that is essential to bone. Strontium is a trace mineral that may help contribute to bone integrity, increasing BMD when bone anabolic processes are impaired by hormonal imbalance, aging, genes, or behaviors such as smoking, drinking carbonated beverages, or excessive alcohol intake. Strontium is abundant in nature and occurs naturally in sea water and in soil. Good plant sources include cabbage, parsley, grapefruit, nuts, asparagus, onion, carrot, tomato, dandelion, oranges, and cucumber. Though not considered an essential nutrient, early research suggested that strontium levels in water may be inversely related to the incidence of dental caries and that it may help control processes of bone resorption.¹

Bone and Skeletal Health†

A growing body of research highlights the benefits of strontium in skeletal health. A head-to-head study comparing SrC to strontium ranelate (SrR) in rats found both forms deliver equivalent amounts of elemental strontium. In the study, equivalent doses of both forms of strontium or placebo were given and assimilation into bone was compared for 19 weeks and followed for another eight weeks. The study confirmed the equivalent delivery of SrC into bone compared to SrR and potential of SrC to support bone structure.²

Recent case studies demonstrate SrC adequately supports BMD in older women. The *Journal of Nutritional Health & Food Science* reported three cases of older women who took SrC on their own and saw bone density supported. In these cases, the subjects chose to discontinue traditional therapies.³ Another unique case was published of a woman who decided to take SrC and allow researchers at McMaster University to follow her bone strontium levels for nearly three years. The subject was 68 years old and had no history of supplementing strontium in any form prior to the study. The patient supplemented with two doses of SrC, which provided 680 mg of strontium per day. Researchers measured the changes in strontium levels in her finger and ankle bones. Not only did her bone strontium levels statistically increase after just five days, but it continued to increase until it reached a plateau after one year.⁴ A 2014 clinical trial published in *Bone* recruited 10 female volunteers who also self-supplemented with 680 mg of strontium per day and found similar results upon evaluation.⁵

Two recent clinical studies also demonstrate that SrC supports BMD and bone quality biomarkers. The COMB Study offered

female patients in need of bone support a micronutrient protocol; these patients had refused or failed to respond to past treatments. After 12 months of consecutive supplemental micronutrient therapy with a combination that included vitamin D3, vitamin K2, SrC, magnesium and docosahexaenoic acid (DHA), repeat bone densitometry was performed. The results showed that bone was supported in the hip, spine, and femoral neck sites. Benefits were found in the group taking the micronutrient protocol.⁶ The MOTS Study was a one-year, double-blind, placebo-controlled trial that assessed the effects of nightly melatonin, SrC, vitamin D3 and vitamin K2 (MK-7) (MSDK) on BMD and quality of life in women ages 49-75. Compared to placebo, MSDK supported BMD in the lumbar spine and femoral neck and supported bone biomarkers. This demonstrates improved bone quality and collagen preservation. Quality of life scores, osteoblast activity, and osteoclast activity were also found to be supported.⁶

Strontium Safety†

Past risk analysis studies of SrR have suggested precaution. A recent 2014 review in *Expert Opinion on Drug Safety* concluded good tolerability and safety profiles in long-term studies.^{8,9} Further, the safety and side effects reported in the PREVOS, STRATOS, TROPOS and SOTI trials do not indicate any adverse events with long-term strontium use.

Directions

1 capsule two times per day or as recommended by your health care professional. For maximum absorption, do not consume with calcium supplements.

Does Not Contain

Gluten, corn, yeast, artificial colors and flavors.

Cautions

If you are pregnant or nursing, consult your physician before taking this product.

Supplement Facts ^{v2}		
Serving Size 1 Capsule		
Servings Per Container 60		
	Amount Per Serving	% Daily Value
Strontium (from 950 mg of Strontium Citrate)	300 mg	*
* Daily Value not established		

References

1. Shuang Tan, Binbin Zhang, Xiaomei Zhu, Ping Ao, Huajie Guo, Weihong Yi, Guang-Qian Zhou, "Deregulation of Bone Forming Cells in Bone Diseases and Anabolic Effects of Strontium-Containing Agents and Biomaterials", *BioMed Research International*, vol. 2014, Article ID 814057, 12 pages, 2014.
2. Wohl GR, Chettle DR, Pejović-Milić A, et al. Accumulation of bone strontium measured by in vivo XRF in rats supplemented with strontium citrate and strontium ranelate. *Bone*. 2013;52(1):63-69.
3. Mirza FS, Azim S, Bhargava A (2016) Change in Bone Mineral Density with Strontium Citrate: An Illusion or Reality. *J Nutrition Health Food Sci* 4(3): 1-3.
4. Moise H, Adachi JD, Chettle DR, Pejović-Milić A. Monitoring bone strontium levels of an osteoporotic subject due to self-administration of strontium citrate with a novel diagnostic tool, in vivo XRF: a case study. *Bone*. 2012;51(1):93-97.
5. Moise H, Chettle DR, Pejović-Milić A. Monitoring bone strontium intake in osteoporotic females self-supplementing with strontium citrate with a novel in-vivo X-ray fluorescence based diagnostic tool. *Bone*. 2014;61:48-54.
6. Stephen J. Genuis, Thomas P. Bouchard, "Combination of Micronutrients for Bone (COMB) Study: Bone Density after Micronutrient Intervention", *Journal of Environmental and Public Health*, vol. 2012, Article ID 354151, 10 pages, 2012.
7. Maria S, Swanson MH, Enderby LT, et al. Melatonin-micronutrients Osteopenia Treatment Study (MOTS): a translational study assessing melatonin, strontium (citrate), vitamin D3 and vitamin K2 (MK7) on bone density, bone marker turnover and health related quality of life in postmenopausal osteopenic women following a one-year double-blind RCT and on osteoblast-osteoclast co-cultures. *Aging (Albany NY)*. 2017;9(1):256-285.
8. Reginster JY. Cardiac concerns associated with strontium ranelate. *Expert Opin Drug Saf*. 2014;13(9):1209-1213.
9. Cianferotti L, D'Asta F, Brandi ML. A review on strontium ranelate long-term antifracture efficacy in the treatment of postmenopausal osteoporosis. *Ther Adv Musculoskelet Dis*. 2013;5(3):127-139.
10. Audran M, Jakob FJ, Palacios S, et al. A large prospective European cohort study of patients treated with strontium ranelate and followed up over 3 years. *Rheumatol Int*. 2013;33(9):2231-2239.