

Energy metabolism and neurological health[‡]

B-Complex

DESCRIPTION

B-Complex is a combination of B vitamins, including vitamins B1, B2, B3, B5, B6, B12, biotin and folate (as Metafolin L-5-MTHF), all of which are provided in their optimal bioavailable and functional forms.[‡]

INDICATIONS

- Support for energy metabolism[‡]
- Support for neurological health[‡]

FUNCTIONS AND MECHANISM OF ACTION

B vitamins play important roles in nearly all physiological systems in the body. Some of the key supportive roles involve nutrient and energy metabolism. B vitamins are essential to basal mitochondrial function as metabolic coenzymes, playing a key role in energy production and nutrient metabolism. Optimal levels of B vitamins have been associated with cellular, cardiovascular, neurological and psychological health. Methylation is a key regulator of genetic expression and cellular function. Healthy methylation is vital for cellular health, DNA synthesis and energy metabolism. At the molecular level, the main function of folate is to donate methyl groups in key biochemical reactions occurring in blood cells, neurons, the vasculature and many other tissues. In addition, folate promotes the production of healthy DNA. In conjunction with vitamins B12 and B6, folate helps to support healthy homocysteine metabolism, cardiovascular function and neurological health. Folate is offered in this formula as 5-methyltetrahydrofolate (5-MTHF), the universally metabolized and biologically active form of folate, providing support for individuals with genetic variations in the MTHFR gene. Genetic variations affecting activation and metabolism of supplemental folic acid are highly prevalent. Through bypassing several enzymatic activation steps, 5-MTHF is directly usable by the body and provides all of the benefits of folic acid regardless of functional or genetic variations. Vitamin B12, as methylcobalamin, works with 5-MTHF to support methylation of homocysteine. Methylcobalamin also supports neuronal function and healthy brain cell activity.[‡]

SUGGESTED USE

As a dietary supplement, adults take 1 capsule, 1-2 times daily, as directed by your health professional.

WARNING

If you are pregnant, nursing, have any health condition or taking any medication, consult your health professional before using this product.

Keep out of reach of children.

STORAGE

Store in a cool, dry place, away from direct light. Use only if safety seal is intact.

FORMULA (#202676)

Supplement Facts		
Serving Size	1 Vegetarian Capsule	
Servings Per Container 60		
	Amount Per Serving	%DV
Thiamine [vitamin B ₁ , as thiamine HCl]	100 mg	8,333%
Riboflavin [vitamin B ₂ and 60% riboflavin 5' phosphate]	12.7 mg	977%
Niacin [vitamin B ₃ , as niacinamide] and inositol hexaniacinate]	108 mg	675%
Vitamin B ₆ [as pyridoxine HCl and 40% pyridoxal 5' phosphate]	16.7 mg	982%
Folate [as Metafolin [®] , L-5-MTHF] (400 mcg L-5-MTHF)	667 mcg DFE	167%
Vitamin B ₁₂ [as methylcobalamin]	400 mcg	16,667%
Biotin.....	400 mcg	1,333%
Pantothenic Acid [vitamin B ₅ , as calcium pantothenate]	50 mg	1,000%

%Daily Value (DV)

Other ingredients: Hydroxypropyl methylcellulose (capsule), ascorbyl palmitate

Gluten-free, Non-GMO

Metafolin is a registered trademark of Merck KGaA, Darmstadt, Germany.

REFERENCES

1. Huskisson E, Maggini S, Ruf M. *J Int Med Res.* 2007 May-Jun;35(3):277-89.
2. Depeint F, Bruce WR, Shangari N. *Chemico-Biological Interactions.* 2006; 123; 94–112.
3. Lonsdale D. *Evid Based Complement Alternat Med.* 2006 Mar; 3(1): 49–59.
4. O'Connell BS. *Diabetes Spectrum.* 2001; 14(3). 133-148.
5. Revilla-Monsalve C, Zendejas-Ruiz I, Islas-Andrade S, et al. *Biomed Pharmacother.* 2006 May;60(4):182-5.
6. Maebashi M, Makino Y, Furukawa Y, et al. *J Clin Biochem Nutr.* 1993; 14(3); 211-218.
6. Olpin SE, Bates CJ. *Br J Nutr* 1982;47:577-88
7. Xiang X, Liu Y, Zhang X, et al. *Wei Sheng Yan Jiu.* 2015 Mar;44(2):185-9, 195.
8. Dakshinamurki K, Mistry SP. *J Biol Chem.* 1963 Jan;238:297-301.
9. Foulon V, Snijders M, Huysmans E, et al. *J Biol Chem.* 2005; 280. 9802-9812.
10. Glier MB, Green TJ, Devlin AM. *Mol Nutr Food Res* 2014 Jan;58(1):172-82.
11. Smulders YM, Smith DE, Kok RM, et al. *Br J Haematol.* 2006 Mar;132(5):623-9.
12. Brunaud L, Alberto JM, Ayav A, et al. *Clin Chem Lab Med.* 2003 Aug;41(8):1012-9.
13. Porter K, Hoey L, Hughes CF, Ward M, McNulty H. *Nutrients.* 2016 Nov 16;8(11). pii: E725.
14. Terry MB, Delgado-Cruzata L, Vin-Raviv N, et al. *Epigenetics.* 2011; 6(7); 828-37.
15. Koury MJ, Ponka P. *Annu Rev Nutr.* 2004;24:105-31.
16. Cohick PL, Bhattacharjee M. *Eur J Clin Nutr* 2011;65, 423–424.
17. Combs GF. *Health and Fitness.* pp. 342.
18. Merete C, Falcon LM, Tucker KL. *J Am Coll Nutr.* 2008 Jun; 27(3): 421–427.
19. Rose DP. *J Clin Pathol.* 1972 Jan; 25(1): 17–25.
20. Stahl SM. *J Clin Psychiatry.* 2008 Sep;69(9):1352-3.
21. Crivello NA, Blusztajn JK, Joseph JA, et al. *Nutr Res.* 2010 Oct;30(10):722-30.
22. Miller AL. *Altern Med Rev.* 2008 Sep;13(3):216-26.
23. Fenech M. *Adv Nutr.* 2017 Nov 15;8(6):958-970.
24. Skipper HE, Mitchell JH, Bennett, LL. *Cancer Res.* 1950. 10;510.
25. Antoniades C, Shirodaria C, Warrick N, et al. *Circulation.* 2006 Sep 12;114(11):1193-201.
26. Cagnacci A, Canoletta M, Volpe A. *Eur J Clin Nutr.* 2009 Oct;63(10):1266-8.
27. Robinson K, Arheart K, Refsum H, et al. *Circulation.* 1998; 97: 437-443.
28. Cui R, Iso H, Date C, et al. *Stroke.* 2010 Jun;41(6):1285-9.
29. Schnyder G, Roffi M, Flammer Y, et al. *JAMA.* 2002 Aug 28;288(8):973-9.
30. MacKenzie KE, Wiltshire EJ, Gent R, et al. *Pediatrics.* 2006 Jul;118(1):242-53.
31. Almeida OP, Marsh K, Alfonso H, et al. *Ann Neurol.* 2010 Oct;68(4):503-10.
32. Tanaka T, Scheet P, Biusti B, et al. *Amer J Hum Genetics* Apr 2009. 84: 477–482.
33. Douaud G, Refsum H, de Jager CA, et al. *Proc Natl Acad Sci U S A.* 2013 Jun 4;110(23):9523-8.
34. den Heijer M, Brouwer IA, Bos GM, et al. *Arterioscler Thromb Vasc Biol.* 1998 Mar;18(3):356-61.
35. Iamopas O, Ratanachu-ek S, Chomtho S. *J Med Assoc Thai.* 2014 Jun;97 Suppl 6:S195-204.
36. Loria-Kohen V, Gmez-Candela C, Palma-Milla S, et al. *Nutr Hosp.* 2013 May-Jun;28(3):807-15.
37. Douaud G, Refsum H, de Jager CA, et al. *Proc Natl Acad Sci U S A* 2013;110(23):9523-8
38. Jacobs AM, Cheng D. *Rev Neurol Dis.* 2011;8(1-2):39-47.
39. Lvblad K, Ramelli G, Remonda L, et al. *Pediatr Radiol.* 1997 Feb;27(2):155-8.
40. K, Ramelli G, Remonda L, et al. *Pediatr Radiol.* 1997 Feb;27(2):155-8.
41. Douaud G, Refsum H, de Jager CA, et al. *Proc Natl Acad Sci U S A.* 2013 Jun 4;110(23):9523-8.
42. Ryan-Harshman M, Aldoori W. *Can Fam Physician.* 2007 Jul; 53(7): 1161–1162.
44. van Asselt DZ, Pasman JW, van Lier HJ, et al. *J Gerontol A Biol Sci Med Sci.* 2001 Dec;56(12):M775-9.
45. Yaqub BA, Siddique A, Sulimani R. *Clin Neurol Neurosurg.* 1992;94(2):105-11.
46. Loria-Kohen V, Gmez-Candela C, Palma-Milla S, et al. *Nutr Hosp.* 2013 May-Jun;28(3):807-15.
47. Jacobs AM, Cheng D. *Rev Neurol Dis.* 2011;8(1-2):39-47.
48. Kienecker EW, Becker KW, Dick P. *Klin Wochenschr.* 1990 Jan 19;68(2):146-8.

49. Hasegawa K, Mikuni N, Sakai Y. *Nihon Yakurigaku Zasshi*. 1978 Sep;74(6):721-34.
50. Fujii A, Matsumoto H, Yamamoto H. *Gen Pharmacol*. 1996 Sep;27(6):995-1000.
51. Zhang X, Chen S, Li L, et al. *Neuropharmacol*. 2008 Jun;54(7):1112-9.
52. Lamers Y, Prinz-Langenohl R, Br mswig S, Pietrzik K. *Am J Clin Nutr*. 2006 Jul;84(1):156-61.
53. Prinz-Langenohl R, Br mswig S, Tobolski O, et al. *Br J Pharmacol*. 2009 Dec; 158(8): 2014–2021.
54. Seremak-Mrozikiewicz A. *Ginekol Pol*. 2013 Jul;84(7):641-6.
55. Hung KL, Wang CC, Huang CY, Wang SJ. *Eur J Pharmacol*. 2009 Jan 14;602(2-3):230-7.
56. Lin Y1, Desbois A, Jiang S, Hou ST. *Neuroreport*. 2004 Oct 5;15(14):2241-4.
57. Akaike A, Tamura Y, Sato Y, Yokota T. *Eur J Pharmacol*. 1993 Sep 7;241(1):1-6.
58. Kikuchi M, Kashii S, Honda Y, et al. *Invest Ophthalmol Vis Sci*. 1997 Apr;38(5):848-54.
59. Obeid R, Herrmann W. *FEBS Lett*. 2006 May 29;580(13):2994-3005.
60. Sachser RM, Haubrich J, Lunardi PS, de Oliveira Alvares L. *Neuropharmacol*. 2017 Jan;112(Pt A):94-103.